

Contribution of Type Ia and Type II Supernovae for Intra-Cluster Medium Enrichment

Shigehiro Nagataki¹ and Katsuhiko Sato^{1,2,3}

¹Department of Physics, School of Science, the University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113, Japan

²Research Center for the Early Universe, School of Science, the University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113, Japan

³ The Institute of Physical and Chemical Research, Wako, Saitama, 351-01, JAPAN

Received _____; accepted _____

ABSTRACT

The origin of the chemical composition of the intracluster medium (ICM) is discussed in this paper. In particular, the contribution from Type Ia supernovae (SNe Ia) to the ICM enrichment is shown to exist by adopting the fitting formulas which have been used in the analysis of the solar system abundances. Our analysis means that we can use the frequency of SNe Ia relative to SNe II as the better measure than $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ for estimating the contribution of SNe Ia. Moreover, the chemical compositions of ICMs are shown to be similar to that of the solar system abundances. We can also reproduce the sulfur/iron abundance ratio within a factor of 2, which means that the abundance problem of sulfur needs not to be emphasized too strongly. We need more precise observations to conclude whether ICMs really suffer the shortage problem of sulfur or not.

Subject headings: intracluster medium (ICM): individual (Abell 496, Abell 1060, Abell 2199, AWM 7)—X-rays—Supernovae

1. Introduction

Primary work on intracluster medium (ICM) enrichment has focused on iron, since this was the only element that was accurately measured in a large number of clusters, prior to the launch of *Advanced Satellite for Cosmology and Astrophysics* (*ASCA*, Tanaka et al. 1994). However, *ASCA* X-ray observations of clusters of galaxies provide the first opportunity to determine the element abundances of O, Ne, Mg, Si, S, Ca, Ar, and Fe for the bulk of the intracluster medium (ICM) in clusters of galaxies (Mushotzky 1995). This abundance ratio pattern in the ICM provided a unique tool to probe the origin of these heavy elements. In particular, a precise analysis of these elemental abundances was done for four brightest, moderate temperature clusters of galaxies (Abell 496, 1060, 2199, and AWM 7). The mean abundances of O, Ne, Si, S and Fe were reported as 0.48, 0.62, 0.65, 0.25, and 0.32, respectively, relative to the solar photospheric abundances (Mushotzky et al. 1996).

Loewenstein & Mushotzky (1996) tried to explain the observed abundance ratios of these elements with respect to Fe (since Fe is the most accurately measured element) by the calculated yields from Type II supernovae (SNe II). They concluded that the abundance ratio pattern of ICM is very similar to that of SNe II. However, they emphasized that a significant SNe Ia contribution to the Fe enrichment could not be ruled out. In fact, Ishimaru & Arimoto (1997) concluded that a SNe Ia iron contribution of 50% or higher to the ICM enrichment could not be ruled out and might be favored by the observations. In the end, Gibson et al. (1997) pointed out the large uncertainties of the adopted massive star physics and uncertainties in the resulting SNe II’s chemical composition, which make it impossible to determine the contribution of SNe Ia to the iron enrichment of ICM. As a result, we have no way of estimating the contribution of SNe Ia to the ICMs.

There is another problem pointed out by Mushotzky et al. (1996), Loewenstein &

Mushotzky (1996), and Gibson et al. (1997). It is the lack of sulfur in the ICMs. Gibson et al. (1997) insist that there would appear to be no escape from the fact that the sulfur abundances are at odds with a SNe II-dominated origin to the ICM iron abundance (although the uncertainties are large). This appears to be entirely independent of adopted SNe II yields (Gibson et al. 1997). Quantitatively, the observed ratio S/Fe favors the ICM SNe Ia iron fractional contribution to be larger than 80%. On the other hand, O/Fe, Si/Fe, Mg/Fe, and Ne/Fe favor a ratio under 60%.

In this paper, we adopt the fitting formula used in the analysis of the solar system abundances (Tsujimoto et al. 1995) and a χ^2 fitting formula to obtain the relative contributions of SNe Ia and SNe II to the enrichment of ICM. We anticipate that these formulas will become good measures for estimating the fractional contribution of SNe Ia. We will also adopt many models of SNe Ia (model W7 and WDD2, Nomoto et al. 1997) and SNe II (Nagataki et al. 1997) which take into account the effect of axisymmetric explosion, in order to reexamine the shortage problem of sulfur.

In section 2, the results of X-ray observations of clusters of galaxies are presented. In section 3, we show a range of calculated yields from SNe Ia and SNe II. In section 4, we present our analysis and results. Summary and discussion are given in section 5.

2. Observations

We show the abundances derived from SIS (solid state imaging spectrometer) in Table 1. $[M_i/Fe]$ is defined as $\log(M_i/Fe)_{\text{ICM}} - \log(M_i/Fe)_{\text{solar}}$. The error bars indicate 90% confidence intervals. We note that the meteoritic abundance scale of Anders & Grevesse (1989) is adopted. Oxygen, silicon, and iron have the best-determined abundances. On the

other hand, the accuracy for magnesium, argon, and calcium is very poor (Mushotzky et al. 1996).

EDITOR: PLACE TABLE 1 HERE.

3. Models of SNe Ia and SNe II

There is a range of SNe II models available in the literature. In the present situation, there are large differences in the predicted yields since there is a wide variety of input physics (e.g., criterion for convection, reaction rates, and the way to initiate the shock wave) (Aufderheide et al. 1991; Gibson et al. 1997).

In order to compare the differences quantitatively, we make use of the formalism presented by Ishimaru & Arimoto (1997). The i th element yield averaged over the SNe II progenitor initial mass function (IMF), $\phi \propto m^{-x}$, is written as

$$\langle y_{i,\text{SNII}} \rangle = \frac{\int_{m_l}^{m_u} y_{i,\text{SNII}}(m) \phi(m) m^{-1} dm}{\int_{m_l}^{m_u} \phi(m) m^{-1} dm}.$$

The bounds for SNe II progenitors are taken to be $m_l = 10M_\odot$ and $m_u = 50M_\odot$, respectively, and an IMF slope $x = 1.35$ (Salpeter 1955) is adopted throughout. While this slope is of prime importance for arguments concerning the absolute mass of elements in the ICM (e.g., Loewenstein & Mushotzky 1996), it is less so for abundance ratios (Gibson et al. 1997).

In imitation of Gibson et al. (1997), we summarize $\langle y_{i,\text{SNII}} \rangle$ for a range of SNe II models in Table 2. We add the yields predicted by Nagataki et al. (1997) and explain it here. They performed 2-dimensional hydrodynamical calculations and studied the changes

of the chemical compositions using a large nuclear reaction network containing 242 nuclear species. Model N97S1 in Table 2 is a spherical explosion model, N97A1 and N97A3 are models of axisymmetric explosions in an increasing order of the degree of deviation from spherical symmetry. For the axisymmetric models N97A1 and N97A3, the initial velocity behind the shock wave is assumed to be radial and proportional to $r \times \frac{1+\alpha \cos(2\theta)}{1+\alpha}$, where r , θ , and α are the radius, the zenith angle and the model parameter that determines the degree of deviation from spherical symmetry, respectively. Nagataki et al. (1997) took $\alpha = 0$ for model N97S1, $\alpha = \frac{1}{3}$ for model N97A1, and $\alpha = \frac{7}{9}$ for model N97A3. The larger α gets, the more asymmetric the explosion becomes. They used the same distribution also for the thermal energy. Half of the total energy appears as kinetic energy and the other half as thermal energy.

As we can see from Table 2, almost all $\langle y_{i,\text{SNII}} \rangle$ of Nagataki et al. (1997) are in the range of uncertainties of theoretical predictions. However, the amount of sulfur tends to become smaller as α gets larger. In fact, N97A3 predicts the least amount of sulfur among all models. This tendency is expected to be good for the reproduction of the observed amount of sulfur in ICMs.

We also show the SNe Ia yields $\langle y_{i,\text{SNeIa}} \rangle$ at the lower portion of Table 2. W7 is the model of the simple deflagration and WDD2 is that of the delayed detonation (Nomoto et al. 1997). As pointed by Gibson et al. (1997), all SNe Ia yields have been tied exclusively to the W7 model. In this paper, the uncertainty of SNe Ia yields is taken into consideration for the first time.

EDITOR: PLACE TABLE 2 HERE.

4. Analysis & Results

4.1. Previous Analysis

At first, the analysis which is done in the previous papers is reexamined. We calculate the abundances ratios $[M_i/\text{Fe}]$ for $M_i = \text{O}, \text{Ne}, \text{Si}, \text{and S}$ as a function of $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$. $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ is the contribution of SNe Ia to the ICM iron enrichment and is given as

$$\frac{M_{\text{Fe,SNIa}}}{M_{\text{Fe,total}}} = \frac{r < y_{\text{Fe,SNIa}} >}{r < y_{\text{Fe,SNIa}} > + (1-r) < y_{\text{Fe,SNII}} >}, \quad (0 \leq r \leq 1) \quad (1)$$

where r indicates the relative occurrence frequency of SNe Ia. The result is shown in Figure 1. Horizontal dot lines mean the average of *ASCA* SIS data.

EDITOR: PLACE FIGURE 1 HERE.

$[\text{O}/\text{Fe}]$ and $[\text{Ne}/\text{Fe}]$ are not changed even when the model of SNe II is changed. This is because the abundances of O and Ne are not determined mainly by the explosive nucleosynthesis but by the nucleosynthesis during the stellar evolution. We also note that the abundance of Fe is mainly determined by the amount of ^{56}Ni , which decays to ^{56}Fe . Since the position of the mass cut is also determined by the amount of ^{56}Ni , Fe abundance is not changed among three models, too (Nagataki et al. 1997). On the other hand, $[\text{Si}/\text{Fe}]$ and $[\text{S}/\text{Fe}]$ are sensitive to the selection of the model of SNe II. This is because Si and S are mainly synthesized during the explosive nucleosynthesis. Si and S abundances tend to decrease along with the degree of deviation from spherical symmetry. Because of this, the $[\text{S}/\text{Fe}]$ ratios give smaller $M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}}$ when W7 and the axisymmetric models are adopted.

As a result, the observations of O, Ne, Si, S, and Fe favor the ICM SNe Ia iron fractional contribution in the range $30\% \leq M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}} \leq 60\%$ for the models W7

and N97A3. On the other hand, it is in the range $30\% \leq M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}} \leq 90\%$ for the models W7 and N97S1. Even worse, the averaged [S/Fe] ratio can not be reproduced when model WDD2 is adopted, although the theoretical [S/Fe] ratio with a proper ratio of $M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}}$ can be within the error bars.

As we see, the ratio of $M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}}$ is very hard to determine, which was also indicated in the previous papers (Loewenstein & Mushotzky 1996; Ishimaru & Arimoto 1997; Gibson et al. 1997).

4.2. $g(\zeta)$ and χ^2 – fitting

Next, we define y_{SNIa} as the sum of $\langle y_{i,\text{SNIa}} \rangle$, which is the heavy element mass of SNe Ia. y_{SNIa} is defined in the same way. We also define the abundance pattern x_i as

$$x_i = \zeta y_{i,\text{SNIa}}/y_{\text{SNIa}} + (1 - \zeta)y_{i,\text{SNIa}}/y_{\text{SNIa}} \quad (0 \leq \zeta \leq 1) \quad (2)$$

which is to be compared with the observed $x_{i,\text{ICM}}$. $x_{i,\text{ICM}}$ is defined as $Z_i/\sum_i Z_i$, where Z_i is the observed abundance of the i -th element per unit mass. ζ is the mass fraction of SNe Ia's matter in ICM. The relation between r and ζ is written as

$$\frac{r}{1-r} = \frac{\omega_{\text{II}} y_{\text{II}}}{\omega_{\text{Ia}} y_{\text{Ia}}} \frac{\zeta}{(1-\zeta)} \quad (3)$$

where ω_{Ia} and ω_{II} represent the mass fraction of heavy elements ejected into the interstellar gas from SNe Ia and SNe II, respectively. These values are estimated to be 0.27 and 0.22 in the solar neighborhood from the numerical calculation (Tsujimoto et al. 1995).

The most probable value of $\zeta = \zeta_p$ is determined by minimizing the following function (Yanagida et al. 1990):

$$g(\zeta) = \sum_{i=1}^n [\log x_{i,\text{ICM}} - \log x_i]^2/n \quad (4)$$

where i runs over the heavy elements considered in the minimization procedure. We summarize those elements in Table 1. We will give another method to obtain ζ_p , that is, χ^2 fitting. As is well known,

$$\chi^2 = \sum_{i=1}^n \left(\frac{x_i(\zeta) - x_{i,\text{ICM}}}{\sigma_i} \right)^2 \quad (5)$$

obeys the χ^2 distribution for n degrees of freedom. We note that σ_i is the standard deviation for $x_{i,\text{ICM}}$. Since we have no information about σ_i , we use the error bars which indicate 90% confidence intervals (Ishimaru & Arimoto 1997) instead.

We show the results for $g(\zeta)$ and χ^2 in Figure 2, 3, 4, and 5. It should be noted that $g(\zeta)$ and χ^2 have a local minimum/maximum for each model, which means there is a favorable value of ζ for the reproduction of the chemical composition of ICMs. Moreover, ζ_p is in the range of $0 < \zeta_p < 0.1$ for almost all clusters, which represents that the contributions of SNe Ia to the ICM enrichment are similar among four ICMs. It should also be noted that $\zeta_p = 0.09 \pm 0.01$ is obtained for the solar system abundances (Tsujimoto et al. 1995), which will mean that the chemical compositions of ICMs are similar to that of the solar system abundances.

Now, there is one question: why can ζ_p be determined in the narrow range $\zeta_p \leq 0.1$ in spite of the large uncertainty of the inferred ratio $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$? The answer is shown in Figure 6, which gives the relation between ζ and $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$. As can be seen from the figure, $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ is very sensitive to ζ if ζ is in the range of $0 \leq \zeta_p \leq 0.1$. This is because the mass fraction of Fe in the SNe Ia's ejecta is quite high. As a result, it is impossible to determine $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ exactly, as we showed in section 4.1. On the other hand, we can use ζ as a better measure than $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ to represent the contribution of SNe Ia to the ICM enrichment.

EDITOR: PLACE FIGURE 2 HERE.

EDITOR: PLACE FIGURE 3 HERE.

EDITOR: PLACE FIGURE 4 HERE.

EDITOR: PLACE FIGURE 5 HERE.

EDITOR: PLACE FIGURE 6 HERE.

Finally, Figure 7 and 8 show the normalized abundance pattern $(x_i/x_{\text{Fe}})_{\text{ICM}}/(x_i/x_{\text{Fe}})_{\text{SN}}$ with the most probable value $\zeta = \zeta_p$ which is obtained from the analysis of $g(\zeta)$ for each cluster. The abundance of sulfur seems to be reproduced better by N97A3 than N97S1, to be sure, however, the sulfur abundance can be reproduced within a factor of 2 even by N97S1. We note that the supernova abundances agree with the solar system abundance ratios within a factor of 2-3 for typical species (Hashimoto 1995), which will reflect the precision of the numerical calculations of supernova nucleosynthesis. Because of this, we think the sulfur abundance problem needs not to be emphasized too strongly. We wait for more precise observations to conclude whether ICMs really suffer the shortage problem of sulfur or not.

EDITOR: PLACE FIGURE 7 HERE.

EDITOR: PLACE FIGURE 8 HERE.

5. Summary and Discussion

In this paper, the origin of the chemical composition in ICMs is discussed. At first, the analysis which has been done in the previous papers is reexamined. As expected, the ratio of $M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}}$ is very hard to determine. Quantitatively, the observations of O, Ne, Si, S, and Fe favor the ICM SNe Ia iron fractional contribution in the range $30\% \leq M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}} \leq 60\%$ for the models W7 and N97A3. The range is $30\% \leq M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}} \leq 90\%$ for the models W7 and N97S1. Even worse, the averaged $[\text{S/Fe}]$ ratio can not be reproduced when model WDD2 is adopted. Our analysis shows that $M_{\text{Fe,SNIa}}/M_{\text{Fe,Total}}$ is not a good measure for estimating the contribution of SNe Ia at the moment to the ICMs.

Next, the analysis of $g(\zeta)$ and χ^2 fitting are done in order to investigate whether these can be better measures for SNe Ia's contribution. We have shown that $g(\zeta)$ and χ^2 have a local minimum/maximum for each model, which means there is a favorable value of ζ for the reproduction of the chemical composition of ICMs. Moreover, ζ_p is in the range of $0 < \zeta_p < 0.1$ for almost all clusters, which means that the contributions of SNe Ia to the ICM enrichment are similar to each other and to the solar system abundances. This analysis indicates that ζ is a better measure than $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$.

The reason why ζ_p can be determined in the narrow range $\zeta_p \leq 0.1$ in spite of the large uncertainty of the inferred ratio $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ is that $M_{\text{Fe,SNIa}}/M_{\text{Fe,total}}$ is very sensitive to ζ if ζ is in the range of $[0, 0.1]$. This is because the mass fraction of Fe in the SNe Ia's ejecta is quite high.

Finally, we have investigated the sulfur abundance problem. The normalized abundance pattern $(x_i/x_{\text{Fe}})_{\text{ICM}}/(x_i/x_{\text{Fe}})_{\text{SN}}$ with the most probable value $\zeta = \zeta_p$ which is obtained from

the analysis of $g(\zeta)$ shows that the sulfur abundance can be reproduced within a factor of 2. Taking the uncertainty of the numerical calculation of supernova nucleosynthesis, we think that the sulfur abundance problem needs not to be emphasized too strongly. We need more precise observations to conclude whether ICMs really suffer the shortage problem of sulfur or not. The same applies for Ar and Ca. We hope that there will be a large number of more precise observations of ICMs to have a more advanced discussion.

We would like to thank W. Hillebrandt for useful comments. We are also grateful to V. Saar for his kind review of the manuscript. This research has been supported in part by a Grant-in-Aid for the Center-of-Excellence (COE) Research (07CE2002) and for the Scientific Research Fund (05243103, 07640386, 3730) of the Ministry of Education, Science, and Culture in Japan and by Japan Society for the Promotion of Science Postdoctoral Fellowships for Research Abroad.

REFERENCES

- Anderse E., Grevesse N. 1989, *Geochim. Cosmochin. Acta*, 53, 197
- Supernova and Nucleosynthesis, Princeton: Princeton Univ. Press
- Aufderheide M. B., Baron E., Thielemann F.K. 1991, *ApJ*, 370, 630
- Gibson, B.K., Loewenstein, M., Mushotzky, R. 1997, *astro-ph/9706138*
- Hashimoto M. 1995, *Prog. Theor. Phys.*, 94, 663
- Ishimaru, Y., Arimoto, N. 1997, *PASJ*, 49, 1
- Loewenstein, M., Mushotzky, R. 1996, *ApJ*, 466, 695
- Maeder, A. 1992, *A&A*, 264, 105 (Erratum: 1993, *A&A*, 268, 833)
- Mushotzky, R. 1995, in *New Horizon of X-Ray Astronomy*, ed. F. Makino & T.Ohashi.(Tokyo: Universal Academy), 243
- Mushotzky, R., Loewenstein, M., Arnaud, K.A., Tamura, T., Fukazawa, Y., Matsushita, K. 1996, *ApJ*, 466, 686
- Nagataki, S., Hashimoto, M., Sato, K., and Yamada, S. 1997, *ApJ*, submitted.
- Nomoto, K. et al. 1997b, *astro-ph/9706025*
- Salpeter, E.E. 1955, *ApJ*, 121, 161
- Tanaka, Y., Inoue, H., Holt, S.S. 1994, *PASJ*, 46, L37
- Tsujimoto T., Nomoto K., Yoshii Y., Hashimoto M., Yanagida S., Thielemann F.-K. 1995, *MNRAS*, 277, 945
- Wang, L., Wheeler, J.C., Li, Z.W., Cliechiatti, A. 1996, *ApJ*, 467, 435

Woosley S.E., Weaver T.A. 1995, ApJS, 101, 181

Yanagida, S., Nomoto, K., and Hayakawa, S. 1990, in Proc. 21st International Cosmic Ray Conference, 4, 44




Fig. 1.— Theoretical ratio of relative abundances of heavy elements, $[O/Fe]$, $[Ne/Fe]$, $[Si/Fe]$, and $[S/Fe]$ as a function of the SN Ia fraction in the iron synthesis. Solid, short-dashed, and dot lines correspond to N97S1, N97A1, and N97A3 models, respectively. Left: W7 model is adopted for SNe Ia. Right: WDD2 model is adopted.




Fig. 2.— $g(\zeta)$ as a function of ζ . W7 and WDD2 are used for SNe Ia. Left: Abell 496. Right: Abell 1060.




Fig. 3.— Same as figure 2 but for Abell 2199 and AWM 7.




Fig. 4.— Confidence level (%) of χ^2 fitting as a function of ζ . Left: Abell 496. Right: Abell 1060.




Fig. 5.— Same as figure 4 but for Abell 2199 and AWM 7.




Fig. 6.— Relation between ζ and $M_{\text{Fe,SNeIa}}/M_{\text{Fe,Total}}$. Solid, dot, short-dashed, and long-dashed lines correspond to W7+N97S1, WDD2+N97S1, W7+N97A3, and WDD2+N97A3, respectively.




Fig. 7.— Normalized abundance pattern $(x_i/x_{\text{Fe}})_{\text{ICM}} / (x_i/x_{\text{Fe}})_{\text{SN}}$ with the most probable value $\zeta = \zeta_p$ which is obtained from the analysis of $g(\zeta)$. Left: Abell 496. Right: Abell 1060.




Fig. 8.— Same with figure 7 but for Abell 2199 and AWM 7.

	Abell 496	Abell 1060	Abell 2199	AWM 7
[O/Fe]	+0.15(−0.15, +0.36)	−0.08(−0.33, +0.08)	−0.06(−0.45, +0.15)	−0.01(−0.37, +0.20)
[Ne/Fe]	+0.28(+0.10, +0.43)	+0.15(+0.00, +0.25)	+0.12(−0.12, +0.28)	−0.16(−0.61, +0.07)
[Mg/Fe]	−0.01(−0.43, +0.21)	−0.31(−0.92, −0.06)	−0.05(−0.48, +0.18)	— (< −0.39) [†]
[Si/Fe]	+0.06(−0.09, +0.18)	+0.14(+0.07, +0.21)	+0.26(+0.16, +0.33)	+0.03(−0.08, +0.13)
[S/Fe]	−0.23(−0.62, −0.02)	−0.30(−0.55, −0.15)	−0.31(−0.86, +0.28)	−0.47(−0.97, −0.23)
[Ar [†] /Fe]	−0.18(< +0.20)	−1.31(< −0.23)	— (< −0.31)	— (< −0.52)
[Ca [†] /Fe]	−0.25(< +0.20)	— (< −0.23)	— (< −0.09)	— (< −0.39)
[Ni/Fe]	+0.49(+0.25, +0.65)	+0.23(−0.33, +0.61)	+0.37(+0.00, +0.54)	+0.19(−0.19, +0.40)

Table 1: Abundances of ICM normalized by the meteoritic abundances. [†] : Nuclei omitted in the analysis of $g(\zeta)$.

Yield Source	< $y_{\text{O,SNI}} >$	< $y_{\text{Ne,SNI}} >$	< $y_{\text{Mg,SNI}} >$	< $y_{\text{Si,SNI}} >$	< $y_{\text{S,SNI}} >$	< $y_{\text{Fe,SNI}} >$
A96	0.593	0.101	0.054	n/a	n/a	0.071
T95	1.777	0.232	0.118	0.133	0.040	0.121
T95+M92	0.923	n/a	n/a	n/a	n/a	0.121
W95;A; $10^{-4} Z_{\odot}$	0.806	0.095	0.036	0.104	0.059	0.073
W95;B; $10^{-4} Z_{\odot}$	1.455	0.223	0.066	0.118	0.065	0.085
W95;A; $Z_{\odot}$	1.217	0.181	0.065	0.124	0.058	0.113
W95;B; $Z_{\odot}$	1.664	0.265	0.094	0.143	0.064	0.141
N97S1	1.749	0.229	0.124	0.139	0.042	0.086
N97A1	1.726	0.231	0.126	0.106	0.026	0.086
N97A3	1.694	0.233	0.123	0.098	0.020	0.088
Yield Source	< $y_{\text{O,SNIa}} >$	< $y_{\text{Ne,SNIa}} >$	< $y_{\text{Mg,SNIa}} >$	< $y_{\text{Si,SNIa}} >$	< $y_{\text{S,SNIa}} >$	< $y_{\text{Fe,SNIa}} >$
W7	0.148	0.005	0.009	0.158	0.086	0.744
WDD2	0.069	0.0009	0.005	0.272	0.168	0.700

Table 2: Average stellar yield in solar masses for the SNe grids. A96 = Arnett 1996, T95 = Tsujimoto et al. 1995, M92 = Maeder 1992, W95 = Woosley & Weaver 1995, N97 = Nagataki et al. 1997c. N97S1, N97A1, and N97A3 stand for the model of the spherical explosion and those of the axisymmetric explosion, respectively. The degree of deviation from spherical symmetry is larger in N97A3 than N97A1. W7 is the model of the simple deflagration and WDD2 is that of the delayed detonation (Nomoto et al. (1997)).